Human neuroblastoma cells with acquired resistance to the p53 activator RITA retain functional p53 and sensitivity to other p53 activating agents

Michaelis, M.; Rothweiler, F.; Agha, B.; Barth, S.; Voges, Y.; Loeschmann, N.; von Deimling, A.; Breitling, R.; Doerr, H. Wilhelm; Roedel, F.

Published in:
Cell death & disease

DOI:
10.1038/cddis.2012.35

Suppl. Table 2. p53 status of nutlin-3-adapted sub-lines of the p53 wild-type RITA-adapted UKF-NB-3 sub-line UKF-NB-3^{r}RITA$^{10\mu M}$.

Cell line	p53 status
UKF-NB-3^{r}RITA$^{10\mu M}$	wild-type
UKF-NB-3^{r}RITA$^{10\mu M}$Nutlin$^{10\mu M}$I	N239D (heterozygote)
UKF-NB-3^{r}RITA$^{10\mu M}$Nutlin$^{10\mu M}$II	wild-type
UKF-NB-3^{r}RITA$^{10\mu M}$Nutlin$^{10\mu M}$III	C277F (heterozygote)
UKF-NB-3^{r}RITA$^{10\mu M}$Nutlin$^{10\mu M}$IV	R280I (heterozygote)
UKF-NB-3^{r}RITA$^{10\mu M}$Nutlin$^{10\mu M}$V	wild-type
UKF-NB-3^{r}RITA$^{10\mu M}$Nutlin$^{10\mu M}$VI	C277F (heterozygote)